PAEDIATRIC TUBE FEEDING

PRODUCT FINDER





How old is the paediatric patient?





Infant 0-18 months 0-9 kg



Child 1-6 yr 8-20 kg



Child 7+ yrs 21-45 kg

Is the infant experiencing intolerance to feeding?*



* Feeding intolerance can be identified if ONE symptom of '<2/3 Prescribed daily enteral intake delivered' or 'enteral nutrition (EN) withheld for 48hrs' or 'EN not increased for 48hrs' is present in combination with ONE of 'vomiting and/or diarrhoea' or 'abdominal pain and discomfort' or 'large gastric residue volume'













KEY NUTRITIONAL FEATURES	\
AGE INDICATION	V
INDICATION	V
FORMAT	V
NUTRITIONAL COMPOSITION	V
DIRECTIONS FOR USAGE	V













KEY NUTRITIONAL FEATURES

- 100 kcal/ml
- Optimal energy from protein (10.2%*), with 2.6 g protein /100 kcal
- Intact protein (casein/whey: 40/60)
- Prebiotic fibres scGOS:lcFOS (9:1)
- Nutritionally complete
- With lactose, LCPUFA**, nucleotides,
- Gluten free

* WHO, 2007; Golden MH. Food Nutr Bull. 2009. ** Long Chain Polyunsaturated Fatty Acids













AGE INDICATION ^

• From birth to 18 months old (0-9 kg)













INDICATION ^

Infatrini is an energy and nutrient dense, nutritionally complete feed for infants with (or at risk of) disease related malnutrition and growth faltering, increased energy and nutrient requirements, fluid restrictions and/or limited ability to take oral feed.

Infatrini is a dietary Food for Special Medical Purposes (FSMP) and must be used under medical supervision. It can be used as a sole source of nutrition and is suitable suitable for administration via bottle, naso-enteric tube, gastrostomy or jejunostomy.













FORMAT ^

Infatrini is available in the following presentations:

- 125 ml plastic bottle (liquid)
- 200 ml plastic bottle (liquid)
- 500 ml collapsible bottle (liquid)
- 400 g tin (powder)

















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Infatrini Liquid Per 100kcal/100ml
Energy	kcal	100
Protein	g (en%)	2.6 (10.2)
Carbohydrate	g (en%)	10.2 (40.6)
lactose	g	5.33
polysaccharides	g	4.1
Fat	g (en%)	5.3 (48.1)
linoleic acid (LA)	mg (en%)	701 (6.3)
a-linolenic acid (ALA)	mg	67 (0.6)
arachidonic acid (AA)	mg	25.3
eicosapentaenoic acid (EPA)	mg	6.33
docosahexaenoic acid (DHA)	mg	25.2
Prebiotic scGOS/lcFOS (9:1)	g	0.8
Of which Fibre	g	0.57

Nutrient	Unit	Infatrini Liquid Per 100kcal/100ml
Na	mg	37.0
K	mg	111
Cl	mg	75.0
Ca	mg	100
Р	mg	50.0
Mg	mg	9.00
Fe	mg	1.20
Zn	mg	0.80
Cu	ug	75
Mn	ug	6
Mo	ug	<8
Se	ug	3.75
Cr	ug	<8
	Ug	19.0
Vit A	ug RE	88.0
Vit D	ug	2.40
Vit E (alpha-tocopherol)	mg	2.52
Vit K	ug	6.70
Thiamin	mg	0.15
Riboflavin	mg	0.20
Niacin preformed	mg	0.80
Niacin .	mg NE	1.47
Pant. Acid	mg	0.80
Vit B6	mg	0.11
Folic acid	ug	16.0
Folate	ug ĎFE	26.7
Vit B12	ug	0.30
Biotin	ug	4.00
Vit C	mg	14.0
Choline	mg	31.3
Carnitine	mg	2.00
Taurine	mg	7.00
Myo-inositol	mg	25.0













DIRECTIONS FOR USAGE ^

Infatrini should be used at room temperature. 125 ml bottles can be attached to a Flocare Pack giving set using a Flocare Universal Adaptor. Alternatively, feed can be decanted into a sterile reservoir, taking care to handle aseptically at all times. Maximum hanging time is 4 hours. The collapsible bottle can be attached directly to a Flocare Pack giving set. When directly connected to a giving set (aseptically handled, closed system), the collapsible bottle has a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then the collapsible bottles have a hanging time of max 4 hours.













KEY NUTRITIONAL FEATURES	\vee
AGE INDICATION	V
INDICATION	V
FORMAT	V
NUTRITIONAL COMPOSITION	V
DIRECTIONS FOR USAGE	\













KEY NUTRITIONAL FEATURES

- 1 kcal/ml
- 100% extensively hydrolyzed whey protein
- Optimal energy from protein (10.2%*), with 2.6 g protein /100 kcal
- Nutritionally complete
- With lactose, LCPUFA**, nucleotides, MCT
- Gluten & fibre free

* WHO, 2007; Golden MH. Food Nutr Bull. 2009. ** Long Chain Polyunsaturated Fatty Acids













AGE INDICATION ^

• From birth to 18 months old (0-9 kg)













INDICATION



Infatrini Peptisorb is an energy and nutrient dense, nutritionally complete feed with extensively hydrolysed protein for infants with impaired gastrointestinal (GI) tolerance and with (or at risk of) disease related malnutrition and growth faltering, increased energy and nutrient requirements, fluid restrictions and/or limited ability to take oral feed.

Infatrini Peptisorb is a dietary Food for Special Medical Purposes (FSMP) and must be used under medical supervision. It can be used as a sole source of nutrition and is suitable for administration via bottle, naso-enteric tube, gastrostomy or jejunostomy.













FORMAT

Infatrini Peptisorb is available in the following presentations:

• 200 ml plastic bottle (liquid)













NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Infatrini Peptisorb Per 100kcal/100ml
Energy	kcal	100
Protein	g (en%)	2.6 (10.6)
-	g	-
whey protein	g	2.6
Carbohydrate	g (en%)	10.2 (40.5)
lactose	g	<0.1
polysaccharides	g	7.3
Fat	g (en%)	5.4 (48.9)
linoleic acid (LA)	mg (en%)	686 (6.2)
a-linolenic acid (ALA)	mg	66.4 (0.6)
arachidonic acid (AA)	mg	25.2
eicosapentaenoic acid (EPA)	mg	7.13
- docosahexaenoic acid (DHA)		25.0
-	mg	2.8
Fibre	g	-

Nutrient	Unit	Infatrini Peptisorb Per 100kcal/100ml
Na	mg	37.0
K	mg	111
Cl	mg	75.0
Ca	mg	90.0
P	mg	45.0
Mg	mg	9.00
Fe	mg	1.20
Zn	mg	0.80
Cu	ug	75
Mn	ug	6
Мо	ug	6.00
Se	ug	3.75
Cr	ug	4.00
	ug	19.0
Vit A	ug RE	88.0
Vit D	ug	2.40
Vit E	mg a-TE	2.10
α-tocopherol	mg	2.70
Vit K	ug	6.70
Thiamin	mg	0.15
Riboflavin	mg	0.20
Niacin preformed	mg	0.80
Niacin	mg NE	1.67
Pant. Acid	mg	0.80
Vit B6	mg	0.11
Folic acid	ug	16.0
Folate	ug DFE	26.7
Vit B12	ug	0.30
Biotin	υg	4.00
Vit C	mg	14.0
Choline	mg	32.1
Carnitine	mg	2.00
Taurine	mg	7.00
Myo-inositol	mg	25.0













DIRECTIONS FOR USAGE



Infatrini Peptisorb should be used at room temperature. Before opening, clean bottle top and shoulders with an alcohol swab. Alternatively, feed can be decanted into a sterile reservoir, taking care to handle aseptically at all times. Maximum hanging time is 4 hours.









What are the energy requirements of your paediatric patient?



LOW

e.g. in cases of reduced mobility or immobility, including patients with neurodevelopmental disabilities such as cerebral palsy, congenital neuromuscular disorders, and myelome

STANDARD

In cases of reduced nutritional intake (e.g. due to mechanical or GI dysfunction, congenital abnormalities, reduced appetite)

HIGH

e.g. in cases of patients with fluid restrictions, increased nutrient requirements or increased nutrient losses (e.g. congenital heart disease, disorders of digestion and absorption, cystic fibrosis, burns, critical illness)











KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE













KEY NUTRITIONAL FEATURES



- 0.75 kcal/ml
- Intact protein (casein/whey: 40/60)
- With Multi Fibre
- DHA+EPA
- Lactose & gluten free













AGE INDICATION



• For patients aged 1-6 years or 8-20 kg in body weight













INDICATION



Nutrini Low Energy Multi Fibre has been specially designed for enteral nutritional support for children aged 1-6 years, or 8-20 kg in weight, with (or at risk of) of disease related malnutrition with low energy requirements. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.













FORMAT ^

Nutrini Low Energy Multi Fibre is available in the following presentations

- 200 ml plastic bottles
- 500 ml collapsible bottles

















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini Low Energy MF /100 kcal	Nutrini Low Energy MF /100 ml
Energy	kcal	100	75
Protein	g (en%)	2.75 (11)	2.1 (11)
-	g	1.1	0.8
-	g	1.6	1.2
Carbohydrate	g (en%)	12.4 (49.4)	9.3 (49.4)
Fat	g (en%)	4.4 (39.6)	3.3 (39.6)
Fibre	g	1.0	0.8

Nutrient	Unit	Nutrini Low Energy MF /100 kcal	Nutrini Low Energy MF /100 ml
Na	mg	80	60
K	mg	179	134
Cl	mg	93	70
Ca	mg	80	60
P	mg	67	50
Mg	mg	15	11
Fe	mg	1.3	1.0
Zn	mg	1.3	1.0
Cu	ug	120	90
Mn	mg	0.20	0.15
F	mg	0.09	0.07
Мо	ug	5.3	4.0
Se	ug	4.0	3.0
Cr	ug	4.7	3.5
\(\frac{1}{2}\)	ug ug RE	13	10.0
Vit A Carotenoids	Ug KE	55	41 0.10
Carotenolos	mg ug RE	0.13 12	9.1
Vit D	ug	12 1.3	1.0
Vit E	mg α-TE	1.7	1.3
Vit K	Ug	5.3	4.0
Thiamin	mg	0.20	0.15
Riboflavin	mg	0.21	0.16
Niacin	mg NE	1.5	1.1
Pant. Acid	mg	0.44	0.33
Vit B6	mg	0.16	0.12
Folic acid	ug	20	15
Vit B12	ug	0.23	0.17
Biotin	ug	5.3	4.0
Vit C	mg	13	10.0
Choline	mg	27	20
Carnitine	mg	2.7	2.0
Taurine	mg	10.0	7.5













DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Plastic bottles: can be attached directly to a Flocare bottle giving set.

Alternatively, feed can be decanted into a sterile reservoir, taking care to handle aseptically at all times. Clean bottle top with an alcohol swab before opening. Collapsible bottles: can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.









Is the patient experiencing intolerance to feeding?*



* Feeding intolerance can be identified if ONE symptom of '<2/3 Prescribed daily enteral intake delivered' or 'enteral nutrition (EN) withheld for 48hrs' or 'EN not increased for 48hrs' is present in combination with ONE of 'vomiting and/or diarrhoea' or 'abdominal pain and discomfort' or 'large gastric residue volume'















KEY NUTRITIONAL FEATURES	\
AGE INDICATION	V
INDICATION	V
FORMAT	V
NUTRITIONAL COMPOSITION	V
DIRECTIONS FOR USAGE	V















KEY NUTRITIONAL FEATURES

- 1 kcal/ml
- 100% extensively hydrolyzed whey protein
- 50% MCT, 50% soy oil
- Contains lactose (0.1 g/100 ml)
- Acidic/pasteurized
- Demineralized water
- Fiber free















AGE INDICATION

















INDICATION



Nutrini Peptisorb is a nutritionally complete formula that contains whey-based extensively hydrolyzed protein for the dietary management of disease related malnutrition in combination with malabsorption and/or maldigestion in children 1-6 years old, or 8-20 kg in weight. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.















FORMAT ^

Nutrini Peptisorb is available in the following presentations

• 500 ml collapsible bottles















NUTRITIONAL COMPOSITION ^



Nutrient	Unit	Nutrini Peptisorb / 100 kcal	Nutrini Peptisorb / 100 ml
Energy	kcal	100	100
Protein	g	2.75	2.75
	en%	0,11	O,11
CHO	g	13.7	13.7
	en%	54	54
Sugars	g	0.83	0.83
	g	3.89	3.89
	en%	35	35
	g	1.78	1.78
	g	0	0
Osmolarity	MOsmol/l	295	295

Nutrient	Unit	Nutrini Peptisorb (NIMS)2 / 100 kcal	Nutrini Peptisorb / 100 ml
Na	mg	60	60
K	mg	110	110
Cl	mg	95	95
Ca	mg	60	60
Р	mg	50	50
Mg	mg	11	11
Fe	mg	1.0	1.0
Zn	mg	1.0	1.0
Си	ug	90	90
Mn	mg	0.15	0.15
F	mg	0.07	0.07
Мо	ug	4.0	4.0
Se	ug	3.0	3.0
Cr	ug	3.50	3.50
1	ug ug RE	10	10
Vit A	ug RE	40.9	40.9
Carotenoids	mg ug RE	0.10	0.10
10.5	ug RE	9.1	9.1
Vit D	ug	1.0	1.0
Vit E	mg α-TE	1.25	1.25
Vit K	ug	4.0	4.0
Thiamin	mg	0.15	0.15
Riboflavin	mg_	0.16	0.16
Niacin	mg NE	1.1	1.1
Pant. Acid	mg	0.33	0.33
Vit B6	mg	0.12	0.12
Folic acid	Ug	15	15
Vit B12	Uğ	0.17	0.17
Biotin	Ug	4.00	4.00
Vit C	mg	10.0	10.0
Choline	mg	20	20
Carnitine	mg	2.0	2.0
Taurine	mg	7.5	7.5















DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.











Does the patient require a fibre- or residue-restricted diet?



e.g. in cases of bowel preparation, acute phase of IBD, intestinal failure

NO

YES











Nutrini





KEY NUTRITIONAL FEATURES	\vee
AGE INDICATION	V
INDICATION	V
FORMAT	V
NUTRITIONAL COMPOSITION	\vee
DIRECTIONS FOR USAGE	V













Nutrini





KEY NUTRITIONAL FEATURES

 \wedge

- 1 kcal/ml
- Intact protein (casein/whey: 40/60)
- Lactose free
- DHA+EPA

















AGE INDICATION

• For patients aged 1-6 years or 8-20 kg in body weight

















INDICATION



Nutrini is a nutritionally complete, ready-to-use, fibre free tube feed for the dietary management of disease related malnutrition in children aged 1-6 years old, or 8-20 kg in weight. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.

















FORMAT ^

Nutrini is available in the following presentations

- 500 ml collapsible bottles
- 200 ml plastic bottles





















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini /100 kcal	Nutrini /100 ml
Energy	kcal	100	100
Protein	g	2.8	2.8
	En%	11	11
Carbohydrate	g	12.3	12.3
	En%	49	49
Fat	g	4.4	4.4
	En%	40	40
DHA/EPA	g	0.045	0.045
% of FA	%	1.03%	1.03%
Fibre	g	0	
Osmolality	mOsmol/kg H20	275	275
Osmolarity	mOsmol/l	235	235

Nutrient	Unit	Nutrini /100 kcal	Nutrini /100 ml
Na	mg	60	60
K	mg	110	110
Cl	mg	95	95
Ca	mg	60	60
Р	mg	50	50
Mg	mg	11	11
Fe	mg	1.0	1.0
Zn	mg	1.0	1.0
Cu	ug	81	81
Mn	mg	0.15	0.15
F	mg	0.07	0.07
Мо	ug	4.0	4.0
Se	ug	3.0	3.0
Cr	ug	3.5	3.5
1	ug ug RE	10.0	10.0
Vit A	ug RE	41	41
Carotenoids	mg ug RE	0.10	0.10
		9.1	9.1
Vit D	ug	1.0	1.0
Vit E	mg α-TE	1.3	1.3
Vit K	ug	4.0	4.0
Thiamin	mg	0.15	0.15
Riboflavin	mg_	0.16	0.16
Niacin	mg NE	1.1	1.1
Pant. Acid	mg	0.33	0.33
Vit B6	mg	0.12	0.12
Folic acid	ug	15	15
Vit B12	ug	0.25	0.25
Biotin	ug	4.0	4.0
Vit C	mg	10.0	10.0
Choline	mg	20	20
Carnitine	mg	2.0 7.5	2.0 7.5
Taurine	mg	7.5	7.5

















DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Plastic bottles: can be attached directly to a Flocare bottle giving set. Alternatively, feed can be decanted into a sterile reservoir, taking care to handle aseptically at all times. Clean bottle top with an alcohol swab before opening. Collapsible bottles: can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.

















KEY NUTRITIONAL FEATURES	\vee
AGE INDICATION	V
INDICATION	V
FORMAT	\vee
NUTRITIONAL COMPOSITION	\vee
DIRECTIONS FOR USAGE	V

















KEY NUTRITIONAL FEATURES

 \land

- 1 kcal/ml
- Intact protein (casein/whey: 40/60)
- Lactose free
- DHA+EPA >1.03% of FA
- With Multi Fibre

















AGE INDICATION



• For patients aged 1-6 years or 8-20 kg in body weight

















INDICATION



Nutrini Multi Fibre is a nutritionally complete, ready-to-use, fibre containing tube feed for the dietary management of disease related malnutrition in children 1-6 years old, or 8-20 kg in weight. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.

















FORMAT ^

Nutrini Multi Fibre is available in the following presentations

- 500 ml collapsible bottles
- 200 ml plastic bottles





















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini Multi Fibre /100 ml
Energy	kcal	101
Protein	g	2.7
	En%	11
Carbohydrate	g	12.3
	En%	48
Sugars	g	0.8
Lactose	g	<0.025
Fat	g	4.4
	En%	40
- saturates	g	0.5
- MCT	g	0
- monounsaturates	g	2.6
- polyunsaturates	g	1.3
- eicosapentaenoic acid (EPA)	mg	8,41
- docosahexaenoic acid (DHA)	mg	35.7
Fibre	g	0.8
	En%	1.0

Nutrient	Unit	Nutrini Multi Fibre /100 ml
Na	mg	60
K	mg	110
Čl	mg	95
Ca	mg	60
Р	mg	50
Mg	mg	11
Fe	mg	1.0
Zn	mg	1.0
Cu	mg	0.08
Mn	mg	0.15
F	mg	0.07
Мо	ug	4.0
Se	ug	3.0
Cr	ug	3.5
	ug	10.0
Vit A	ug RE	41
Vit D	ug	1.0
Vit E	mg α-TE	1.3
Vit K	ug	4.0
Thiamin	mg	0.15
Riboflavin	mg_	0.16
Niacin	mg ŇE	1.1
Pant. Acid	mg	0.33
Vit B6	mg	0.12
Folic acid	ug	15
Vit B12	ug	0.25
Biotin	ug	4.0
Vit C	mg	10.0

















DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Plastic bottles: can be attached directly to a Flocare bottle giving set. Alternatively, feed can be decanted into a sterile reservoir, taking care to handle aseptically at all times. Clean bottle top with an alcohol swab before opening. Collapsible bottles: can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.













Is the patient experiencing intolerance to feeding?*



* Feeding intolerance can be identified if ONE symptom of '<2/3 Prescribed daily enteral intake delivered' or 'enteral nutrition (EN) withheld for 48hrs' or 'EN not increased for 48hrs' is present in combination with ONE of 'vomiting and/or diarrhoea' or 'abdominal pain and discomfort' or 'large gastric residue volume'















KEY NUTRITIONAL FEATURES	\
AGE INDICATION	V
INDICATION	V
FORMAT	V
NUTRITIONAL COMPOSITION	V
DIRECTIONS FOR USAGE	V















KEY NUTRITIONAL FEATURES

- 1.5 kcal/ml
- 100% extensively hydrolyzed whey protein
- 53% MCT, 38% soy oil, 9% corn oil
- 0.4 g/100 ml lcFOS/scFOS (1:1)
- Contains lactose (<0.2 g/100 ml)
- Acidic/pasteurized
- Fibre free















AGE INDICATION



• For patients aged 1-12 years















INDICATION



Nutrini Peptisorb Energy is an energy dense, nutritionally complete formula that contains whey-based extensively hydrolyzed protein for the dietary management of disease related malnutrition in combination with malabsorption and/or maldigestion in children (1 year old and onwards).















FORMAT ^

Nutrini Peptisorb Energy is available in the following presentations

• 500 ml collapsible bottles















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini Peptisorb Energy / 100 kcal	Nutrini Peptisorb Energy / 100 ml
Energy	kcal	100	150
Protein	g	2.78	4.2
	En%	11.1	11.1
CHO	g	12.4	18.6
	En%	48.8	48.8
Fat	g	4.39	6.6
	En%	39.5	39.5
	g	2.24	3.4
	g	0.28	0.4
Osmolarity	mOsmol/l	392	392

Nutrient	Unit	Nutrini Peptisorb Energy / 100 kcal	Nutrini Peptisorb Energy / 100 ml
Na	mg	63.2	95
K	mg	112	168
Cl	mg	95.5	143
Ca	mg	86.5	130
Р	mg	66	99
Mg	mg	15.4	23
Fe	mg	1.0	1.5
Zn	mg	1.0	1.5
Си	ug	70	110
Mn	mg	0.10	0.15
F	mg	0.07	0.10
Mo	ug	4.67	7.0
Se	ug	3.14	4.7
Cr	ug	2.5	3.7
	ug_	10.1	15
Vit A	ug RE	41.1	62
Carotenoids	mg ug RE	42.7	64
	ug RE	1.25	1.9
Vit D	ug	1.30	2.0
Vit E	mg α-TE	4.01	6.0
Vit K	ug	0.15	0.23
Thiamin	mg	0.16	0.24
Riboflavin	mg_	1.10	1.7
Niacin	mg NE	0.33	0.50
Pant. Acid	mg	0.12	0.18
Vit B6	mg	18	27
Folic acid	Ug	0.17	0.26
Vit B12	Ug	4.01	6.0
Biotin	Ug	10.0	15
Vit C	mg	25	38
Choline	mg	4.64	7.0
Carnitine	mg	1.97	3.0
Taurine	mg	7.61	11















DIRECTIONS FOR USAGE



Use at room temperature. When connected to the giving set (aseptically handled, closed system) this bottle has a maximum hanging time of 24 hours. If the cap has been removed (used as an open system) then the bottle has a hanging time of maximum 8 hours. Packs can be attached directly to a Flocare Pack giving set.











Does the patient require a fibre- or residue-restricted diet?



e.g. in cases of bowel preparation, acute phase of IBD, intestinal failure

NO YES















KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE

















KEY NUTRITIONAL FEATURES

/\

- 1.5 kcal/ml
- Intact protein (casein/whey: 60/40)
- Fibre free
- Lactose free
- DHA+EPA

















AGE INDICATION

• For patients aged 1-6 years or 8-20 kg in body weight

















INDICATION



Nutrini Energy is a nutritionally complete, energy dense, ready-to-use, fibre free tube feed for the dietary management of disease related malnutrition in children aged 1-6 years old, or 8-20 kg in weight. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.

















FORMAT ^

Nutrini Energy is available in the following presentations

- 500 ml collapsible bottles
- 200 ml plastic bottles





















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini Energy /100 ml	Nutrini Energy /100 kcal
Energy	kcal	150	100
Protein	g	4.0	2.7
	En%	11	11
Carbohydrate	g	18.5	12.3
	En%	49	49
Sugars	g	1.1	0.8
Lactose	g	<0.025	<0.017
Fat	g	6.7	4.4
	En%	40	40
- saturates	g	0.8	0.5
- MCT	g	0	0
- monounsaturates	g	3.9	2.6
- polyunsaturates	g	2.0	1.3
- arachidonic acid (AA)	mg	3.76	2.5
- eicosapentaenoic acid (EPA)	g	12.6	8.43
- docosahexaenoic acid (DHA)	g	53.1	35.4
Fibre	g	0	0

Nutrient	Unit	Nutrini Energy /100 ml	Nutrini Energy /100 kcal
Na	mg	90	60
K	mg	165	110
Cl	mg	143	95
Ca	mg	90	60
Р	mg	75	50
Mg	mg	17	11
Fe	mg	1.5	1.0
Zn	mg	1.5	1.0
Cu	mg	0.12	0.08
Mn	mg	0.23	0.15
F	mg	0.11	0.07
Мо	ug	6.0	4.0
Se	υg	4.5	3.0
Cr	ug	5.3	3.5
I	ug	15	10
Vit A	ug ug RE	61	41
Vit D	ug	1.5	1.0
Vit E	mg α-TE	1.9	1.3
Vit K	ug	6.0	4.0
Thiamin	mg	0.23	0.15
Riboflavin	mg	0.24	0.16
Niacin	mg NE	1.7	1.1
Pant. Acid	mg	0.50	0.33
Vit B6	mg	0.18	0.12
Folic acid	ug	23	15
Vit B12	ug	0.27	0.18
Biotin	ug	6.0	4.0
Vit C	mg	15	10

















DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Plastic bottles: can be attached directly to a Flocare bottle giving set. Alternatively, feed can be decanted into a sterile reservoir, taking care to handle aseptically at all times. Clean bottle top with an alcohol swab before opening. Collapsible bottles: can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.













Is a product including real food ingredients preferred (for e.g. for long-term feeding or optimised tolerance)?

















Nutrini +MIX Multi Fibre





KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE



















KEY NUTRITIONAL FEATURES



- 1.5 kcal/ml
- Intact protein (casein/whey: 57/40)
- With ingredients of fruit and vegetable origin*
- With Multi Fibre
- DHA+FPA
- Lactose free

* 14% of real fruit and vegetable ingredients (carrot 5.5%, zucchini 4.5%, apricot 4%): reconstituted carrot from flakes 5.5%, reconstituted zucchini from flakes 4.5% and reconstituted apricot juice and pulp from powder 4.0%



















AGE INDICATION



• For patients aged 1-17 years old



















INDICATION



Nutrini +MIX Energy Multi Fibre is an energy and nutrient dense, nutritionally complete, ready-to-use, fibre containing liquid formula with real fruit and vegetable ingredients* for children with disease related malnutrition from 1 year onwards in need of enteral tube feeding. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.

* 14% of real fruit and vegetable ingredients (carrot 5.5%, zucchini 4.5%, apricot 4%): reconstituted carrot from flakes 5.5%, reconstituted zucchini from flakes 4.5% and reconstituted apricot juice and pulp from powder 4.0%



















FORMAT ^

Nutrini +MIX Energy Multi Fibre is available in the following presentations

• 500 ml collapsible bottles



















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini +MIX Energy Multi Fibre /100 kcal	Nutrini +MIX Energy Multi Fibre /100 ml
Energy	kcal	100	150
Protein	g (% energy)	3.0 (12)	4.5 (12)
Carbohydrate	g (% energy)	11.5 (45.8)	17.2 (45.8)
Sugars	g	1.3	1.9
Lactose	g	< 0.017	<0.025
Fat	g (% energy)	4.5 (40.1)	6.7 (40.1)
Fibre	g (% energy)	1.1 (2.2)	1.7 (2.2)
Osmolality	mOsml/kg/water	570	570

Nutrient	Unit	Nutrini +MIX Energy Multi Fibre /100 kcal	Nutrini +MIX Energy Multi Fibre /100 ml
Sodium	mg	74.0	111
Potassium	mg	130	195
Chloride	mg	116	174
Calcium	mg	95	143
Phosphorus	mg	63.7	95.5
Magnesium	mg	20	30
Iron	mg	1.41	2.12
Zinc	mg	1.00	1.50
Copper	μg	87	130
Manganese	mg	0.063	0.095
Fluoride	mg	0.07	0.10
Molybdenum	hã	6.0	9.0
Selénium	μg	3.50	5.25
Chromium	þg	3.27	4.9
lodine	μg	13.3	20.0
Vitamin A	pg pg RE	54.5	81.8
Vit D	μg	2.10	3.15 2.34
Vit E	mg αTE	1.56	2.34
Vit K	μg	6.15	9.23
Thiamin	mg	0.15	0.23
Riboflavin	mg	0.14	0.21
Niacin	mg NE 4	1.6	2.4
Pant. Acid	mg	0.67	1.00
Vit B6	mg	0.13	0.20
Folic acid	μg	18.5	27.8
Vit B12	μg	0.22	0.33
Biotin	μg	4.00	6.00
Vit C	mg	10.0	15.0
Choline	mg	30.0	45.0
Inositol	mg	3.36	5.04
Carnitine	mg	4.5	6.75
Taurine	mg	10	15
Carotenoids	mg	0.53	0.8



















DIRECTIONS FOR USAGE



Sedimentation is a normal phenomenon. Check appearance before use and shake well before opening. Do not dilute or add medications to the formula. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.



















KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE



















KEY NUTRITIONAL FEATURES

 \wedge

- 1.5 kcal/ml
- Intact protein (casein/whey: 60/40)
- Lactose free
- DHA+EPA >1.03% of FA
- With Multi Fibre



















AGE INDICATION



• For patients aged 1-6 years or 8-20 kg in body weight



















INDICATION

Nutrini Energy Multi Fibre is a nutritionally complete, energy dense, ready-to-use, fiber containing tube feed for the dietary management of disease related malnutrition in children 1-6 years old, or 8-20 kg in weight. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.



















FORMAT ^

Nutrini Energy Multi Fibre is available in the following presentations

- 500 ml collapsible bottles
- 200 ml plastic bottles























NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini Energy MF /100 kcal	Nutrini Energy MF /100 ml
Energy	kcal	100	150
Protein	g	2,75	4.1
	En%	11	11
Carbohydrate	g	12.3	18.5
	En%	49	49
Fat	g	4.4	6.7
	En%	40	40
DHA/EPA	g	0.044	0.066
% of FA	%	1.04	1.04
Fibre	g	0.5	
Osmolality	mOsmol/kg H20	390	390
Osmolarity	mOsmol/l	300	300

Nutrient	Unit	Nutrini Energy MF /100 kcal	Nutrini Energy MF /100 ml
Na	mg	760	90
K	mg	110	165
Cl	mg	95	143
Ca	mg	60	90
Р	mg	50	75
Mg	mg	11	17
Fe	mg	1.0	1.5
Zn	mg	1.0	1.5
Cu	ug	81	122
Mn	mg	0.15	0.23
F	mg	0.07	0.11
Мо	ug	4.0	6.0
Se	ug	3.0	4.5
Cr	ug	3.5	5.3
I	ug ug RE	10.0	15
Vit A	ug RE	41	61
Carotenoids	mg	0.10	0.15
	ug RE	9.1	13.7
Vit D	ug	1.0	1.5
Vit E	mg α-TE	1.3	1.9
Vit K	ug	4.0	6.0
Thiamin	mg	0.15	0.23
Riboflavin	mg	0.16	0.24
Niacin	mg NE	1.1	1.7
Pant. Acid	mg	0.33	0.5
Vit B6	mg	0.12	0.18
Folic acid	ug	15	23
Vit B12	ug	0.18	0.27
Biotin	ug	4.0	6.0
Vit C	mg	10.0	15
Choline	mg	20	30
Carnitine	mg	2.0	3.0
Taurine	mg	7.5	11.3



















DIRECTIONS FOR USAGE

Shake well before opening. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.













 \wedge



What are the energy requirements of the patient?



STANDARD

In cases of reduced nutritional intake (e.g. due to mechanical or GI dysfunction, congenital abnormalities, reduced appetite)

HIGH

e.g. in cases of patients with fluid restrictions, increased nutrient requirements or increased nutrient losses (e.g. congenital heart disease, disorders of digestion and absorption, cystic fibrosis, burns, critical illness)







Does the patient require a fibre- or residue-restricted diet?



e.g. in cases of bowel preparation, acute phase of IBD, intestinal failure

NO YES













KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE















KEY NUTRITIONAL FEATURES

 \wedge

- 1 kcal/ml
- Intact protein (casein/whey: 40/60)
- Fibre free
- Lactose free
- DHA+EPA















AGE INDICATION



• For patients aged 7-12 years or 21-45 kg in body weight















INDICATION



NutriniMax is a nutritionally complete ready-to-use, fibre free tube feed for the dietary management of disease related malnutrition in children (7-12 years or 21-45 kg). It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.















FORMAT



NutriniMax is available in the following presentations:

• 500 ml collapsible bottles















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	NutriniMax /100 ml
Energy	kcal	100
Protein	g	3.3
	En%	13
Carbohydrate	g	12.3
	En%	49
Polysaccharides	g	11.3
Sugars	g	0.8
Lactose	g	<0.025
Fat	g	4.2
	En%	38
- saturates	g	0.5
- monounsaturates	g	2.4
- polyunsaturates	g	1.3
- eicosapentaenoic acid (EPA)	mg	8.0
- docosahexaenoic acid (DHA)	mg	33.0
Fibre	g	0

Nutrient	Unit	NutriniMax /100 ml
Na	mg	80
K	mg	138
Cl	mg	110
Ca	mg	70
Р	mg	60
Mg	mg	17
Fe	mg	1.3
Zn	mg	1.1
Си	mg	0.11
Mn	mg	0.24
F	mg	0.09
Мо	ug	7.0
Se	ug	4.9
Cr	ug	5.1
	ug	12 61
Vit A	ug RE	
Vit D	ug	0.7
Vit E	mg α-TE	1.3
Vit K	ug	4.5
Thiamin	mg	0.15
Riboflavin	mg	0.16
Niacin	mg NE	1.5
Pant. Acid	mg	0.43
Vit B6	mg	0.15
Folic acid	ug	21
Vit B12	ug	0.24
Biotin	ug	4.0
Vit C	mg	10.0
	1	I /















DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.















KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE

















KEY NUTRITIONAL FEATURES

 \wedge

- 1 kcal/ml
- Intact protein (casein/whey: 40/60)
- Lactose free
- DHA+EPA >1.03% of FA
- With Multi Fibre

















KEY NUTRITIONAL FEATURES

• For patients aged 7-12 years or 21-45 kg in body weight

















INDICATION

NutriniMax Multi Fibre is a nutritionally complete ready-to-use, fibre free tube feed for the dietary management of disease related malnutrition in children (7-12 years or 21-45kg). It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.

















FORMAT ^

NutriniMax Multi Fibre is available in the following presentations:

• 500 ml collapsible bottles

















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	NutriniMax Multi Fibre /100 kcal	NutriniMax Multi Fibre /100 ml
Energy	kcal	100	100
Protein	g	3.3	3.3
	En%	13	13
Carbohydrate	g	12.3	12.3
	En%	49	49
Fat	g	4.2	4.2
	En%	38	38
DHA/EPA	g	0.042	0.042
% of FA	%	1.05	1.05
Fibre	g	1.13	1.13
Osmolarity	mOsmol/l	230	230

Nutrient	Unit	NutriniMax Multi Fibre /100 kcal	NutriniMax Multi Fibre /100 ml
Na	mg	80	80
K	mg	138	138
Cl	mg	110	110
Ca	mg	70	70
P	mg	60	60
Mg	mg	17	17
Fe	mg	1.3	1.3
Zn	mg	1.1	1.1
Си	ug	108	108
Mn	mg	0.24	0.24
F	mg	0.09	0.09
Мо	ug	7.0	7.0
Se	ug	4.9 5.1	4.9
Cr	ug	5.1	5.1
\(\frac{1}{2}\)	ug	11.5	11.5
Vit A Carotenoids	ug RE	61	61
Carotenolos	mg	0.15 14	0.15 14
Vit D	ug ŘE	0.7	0.7
	ug		
Vit E Vit K	mg a-TE	1.3 4.5	1.3 4.5
Thiamin	ug	0.15	0.15
Riboflavin	mg	0.15	0.15
Niacin	mg mg NE	1.5	1.5
Pant. Acid	mg	0.43	0.43
Vit B6	mg	0.15	0.45
Folic acid		21	21
Vit B12	ug ug	0.24	0.24
Biotin	ug	4.0	4.0
Vit C	mg	10.0	10.0
Choline	mg	28	28
Carnitine	mg	3.0	3.0
Taurine	mg	10	10

















DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Maximum hanging time 24 hours. Packs can be attached directly to a Flocare giving set.













Is the patient experiencing intolerance to feeding?*



* Feeding intolerance can be identified if ONE symptom of '<2/3 Prescribed daily enteral intake delivered' or 'enteral nutrition (EN) withheld for 48hrs' or 'EN not increased for 48hrs' is present in combination with ONE of 'vomiting and/or diarrhoea' or 'abdominal pain and discomfort' or 'large gastric residue volume'















KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE















KEY NUTRITIONAL FEATURES

 \wedge

- 1.5 kcal/ml
- 100% extensively hydrolyzed whey protein
- 53% MCT, 38% soy oil, 9% corn oil
- 0.4 g/100 ml lcFOS/scFOS (1:1)
- Contains lactose (<0.2 g/100 ml)
- Acidic/pasteurized
- Fibre free















AGE INDICATION



• For patients aged 1-12 years















INDICATION



Nutrini Peptisorb Energy is an energy dense, nutritionally complete formula that contains whey-based extensively hydrolyzed protein for the dietary management of disease related malnutrition in combination with malabsorption and/or maldigestion in children (1 year old and onwards).















FORMAT ^

Nutrini Peptisorb Energy is available in the following presentations

• 500 ml collapsible bottles















NUTRITIONAL COMPOSITION ^

Nutrient	Unit	Nutrini Peptisorb Energy / 100 kcal	Nutrini Peptisorb Energy / 100 ml
Energy	kcal	100	150
Protein	g	2.78	4.2
	En%	11.1	11.1
CHO	g	12.4	18.6
	En%	48.8	48.8
Fat	g	4.39	6.6
	En%	39.5	39.5
	g	2.24	3.4
	g	0.28	0.4
Osmolarity	mOsmol/l	392	392

Nutrient	Unit	Nutrini Peptisorb Energy / 100 kcal	Nutrini Peptisorb Energy / 100 ml
Na	mg	63.2	95
K	mg	112	168
Cl	mg	95.5	143
Ca	mg	86.5	130
P	mg	66	99
Mg	mg	15.4	23
Fe	mg	1.0	1.5
Zn	mg	1.0	1.5
Cu	ug	70	110
Mn	mg	0.10	0.15
F.	mg	0.07	0.10
Мо	ug	4.67	7.0
Se	ug	3.14	4.7
Cr	ug	2.5	3.7
	ug ug RE	10.1	15
Vit A	ug RE	41.1	62
Carotenoids	mg ug RE	42.7	64
14. 5	ug RE	1.25	1.9
Vit D	ug	1.30	2.0
Vit E	mg α-TE	4.01	6.0
Vit K	ug	0.15	0.23
Thiamin	mg	0.16 1.10	0.24 1.7
Riboflavin	mg mg NE		
Niacin		0.33 0.12	0.50 0.18
Pant. Acid	mg	18	27
Vit B6	mg	0.17	0.26
Folic acid Vit B12	Ug	4.01	6.0
	Ug		15
Biotin Vit C	Ug	10.0 25	38
Choline	mg	4.64	7.0
Carnitine	mg	1.97	3.0
Taurine	mg	7.61	3.0
Taurine	mg	7.01	L " /















DIRECTIONS FOR USAGE

Use at room temperature. When connected to the giving set (aseptically handled, closed system) this bottle has a maximum hanging time of 24 hours. If the cap has been removed (used as an open system) then the bottle has a hanging time of maximum 8 hours. Packs can be attached directly to a Flocare Pack giving set.











Does the patient require a fibre- or residue-restricted diet?



e.g. in cases of bowel preparation, acute phase of IBD, intestinal failure

NO

YES















KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE

















KEY NUTRITIONAL FEATURES

 \wedge

- 1.5 kcal/ml
- Intact protein (casein/whey: 60/40)
- Lactose free
- Fibre free
- DHA+EPA

















AGE INDICATION



• For patients aged 7-12 years or 21-45 kg in body weight

















INDICATION



NutriniMax Energy is an energy dense, nutritionally complete, ready-to-use, fibre free tube feed for the dietary management of disease related malnutrition in children (7-12 years or 21-45 kg). It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.













NutriniMax Energy





FORMAT ^

NutriniMax Multi Fibre is available in the following presentations:

• 500 ml collapsible bottles













NutriniMax Energy





NUTRITIONAL COMPOSITION ^

Nutrient	Unit	NutriniMax Energy /100 kcal	NutriniMax Energy /100 ml
Energy	kcal	100	1100
Protein	g	3.24	4.88
	En%	13	13
Carbohydrate	g	12.3	18.5
	En%	49	49
Fat	g	4.2	6.3
	En%	38	38
DHA/EPA	g	0.042	0.063
% of FA	%	1.05	1.05
Fibre	g	0	0
Osmolality	mOmol/k g H2O	425	425

Nutrient	Unit	NutriniMax Energy /100 kcal	NutriniMax Energy /100 ml
Na	mg	71,2	107
K	mg	124	186
Cl	mg	98	155
Ca	mg	63	95
Р	mg	60	90
Mg	mg	17	26
Fe	mg	1.3	2.0 1.7
Zn	mg	1.3	
Cu	υg	108	162
Mn	mg	0.24	0.36
F	mg	0.09	0.13
Мо	ug	7.0	11
Se	ug	4.9	7.4
Cr	ug	5.1	7.7
l	ug	11.5	17
Vit A	ug ug RE	61.3	92
Carotenoids	mg ug RE	0.15	0.23
	ug RE	14	21 1.1
Vit D	ug	0.7	1.1
Vit E	mg α-TE	1.25	1.9
Vit K	ug	4.5	6.8 0.23
Thiamin	mg	0.15	0.23
Riboflavin	mg mg NE	0.16	0.24 2.23 0.65
Niacin	mg NE	1.5	2.23
Pant. Acid	mg	0.43	0.65
Vit B6	mg	0.15	0.23
Folic acid	ug	21	32
Vit B12	ug	0.23	0.30
Biotin	ug	4.0	6.0
Vit C	mg	10.0	15
Choline	mg	28	43
Carnitine	mg	3.0	4.5
Taurine	mg	10	15













NutriniMax Energy





DIRECTIONS FOR USAGE



Shake well before opening. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.













Is a product including real food ingredients preferred (for e.g. for long-term feeding or optimised tolerance)?



VO

YES

















KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE



















KEY NUTRITIONAL FEATURES

 \wedge

- 1.5 kcal/ml
- Intact protein (casein/whey: 57/40)
- With ingredients of fruit and vegetable origin*
- With Multi Fibre
- DHA+FPA
- Lactose free

* 14% of real fruit and vegetable ingredients (carrot 5.5%, zucchini 4.5%, apricot 4%): reconstituted carrot from flakes 5.5%, reconstituted zucchini from flakes 4.5% and reconstituted apricot juice and pulp from powder 4.0%



















AGE INDICATION



• For patients aged 1-17 years old



















INDICATION



Nutrini +MIX Energy Multi Fibre is an energy and nutrient dense, nutritionally complete, ready-to-use, fibre containing liquid formula with real fruit and vegetable ingredients* for children with disease related malnutrition from 1 year onwards in need of enteral tube feeding. It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.

* 14% of real fruit and vegetable ingredients (carrot 5.5%, zucchini 4.5%, apricot 4%): reconstituted carrot from flakes 5.5%, reconstituted zucchini from flakes 4.5% and reconstituted apricot juice and pulp from powder 4.0%



















FORMAT ^

Nutrini +MIX Energy Multi Fibre is available in the following presentations

• 500 ml collapsible bottles



















NUTRITIONAL COMPOSITION

Nutrient	Unit	Nutrini +MIX Energy Multi Fibre /100 kcal	Nutrini +MIX Energy Multi Fibre /100 ml
Energy	kcal	100	150
Protein	g (% energy)	3.0 (12)	4.5 (12)
Carbohydrate	g (% energy)	11.5 (45.8)	17.2 (45.8)
Sugars	g	1.3	1.9
Lactose	g	< 0.017	<0.025
Fat	g (% energy)	4.5 (40.1)	6.7 (40.1)
Fibre	g (% energy)	1.1 (2.2)	1.7 (2.2)
Osmolality	mOsml/kg/water	570	570

Nutrient	Unit	Nutrini +MIX Energy Multi Fibre /100 kcal	Nutrini +MIX Energy Multi Fibre /100 ml
Sodium	mg	74.0	111
Potassium	mg	130	195
Chloride	mg	116	174
Calcium	mg	95	143
Phosphorus	mg	63.7	95.5
Magnesium	mg	20	30
Iron	mg	1.41	2.12
Zinc	mg	1.00	1.50
Copper	μg	87	130
Manganese	mg	0.063	0.095
Fluoride	mg	0.07	0.10
Molybdenum	hã	6.0	9.0
Selénium	μg	3.50	5.25
Chromium	þg	3.27	4.9
lodine	μg	13.3	20.0
Vitamin A	pg pg RE	54.5	81.8
Vit D	μg	2.10	3.15 2.34
Vit E	mg αTE	1.56	2.34
Vit K	μg	6.15	9.23
Thiamin	mg	0.15	0.23
Riboflavin	mg	0.14	0.21
Niacin	mg NE 4	1.6	2.4
Pant. Acid	mg	0.67	1.00
Vit B6	mg	0.13	0.20
Folic acid	μg	18.5	27.8
Vit B12	μg	0.22	0.33
Biotin	μg	4.00	6.00
Vit C	mg	10.0	15.0
Choline	mg	30.0	45.0
Inositol	mg	3.36	5.04
Carnitine	mg	4.5	6.75
Taurine	mg	10	15
Carotenoids	mg	0.53	0.8



















DIRECTIONS FOR USAGE

Sedimentation is a normal phenomenon. Check appearance before use and shake well before opening. Do not dilute or add medications to the formula. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.









KEY NUTRITIONAL FEATURES

AGE INDICATION

INDICATION

FORMAT

NUTRITIONAL COMPOSITION

DIRECTIONS FOR USAGE



















KEY NUTRITIONAL FEATURES



- 1.5 kcal/ml
- Intact protein (casein/whey: 60/40)
- With Multi Fibre
- Lactose free
- DHA+EPA



















AGE INDICATION



• For patients aged 7-12 years or 21-45 kg in body weight



















INDICATION



NutriniMax Energy Multi Fibre is an energy dense, nutritionally complete, ready-to-use, fiber containing tube feed for the dietary management of disease related malnutrition in children (7-12 years or 21-45 kg). It is suitable for administration via naso-enteric tube, gastrostomy or jejunostomy.



















FORMAT ^

NutriniMax Energy Multi Fibre is available in the following presentations:

• 500 ml collapsible bottles





















Nutrient	Unit	NutriniMax Energy MF/100 kcal	NutriniMax Energy MF/100 r
Energy	kcal	100	150
Protein	g	3.25	4.9
	En%	13	13
Carbohydrate	g	12.3	18.5
	En%	49	49
Fat	g	4.2	6.3
	En%	38	38
DHA/EPA	g	0.042	0.063
% of FA	%	1.05	1.05
Fibre	g	0.75	1.13
Osmolality	mOmol/k g H2O	415	415

Nutrient	Unit	NutriniMax Energy MF /100 kcal	NutriniMax Energy MF /100 ml
Na	mg	71.2	107
K	mg	124	186
Cl	mg	98	147
Ca	mg	63	95
Р	mg	60	90
Mg	mg	17	26
Fe	mg	1.3	2.0
Zn	mg	1.3	1.7
Cu	ug	108	162
Mn	mg	0.24	0.36
F.	mg	0.09	0.13
Mo	ug	7.0	11
Se	ug	4.9	7.4
Cr	ug	5.1	7.7
	ug ug RE	11.5	17
Vit A	ug RE	61.3	92
Carotenoids	mg ug RE	0.15	0.23
Vii D	ug RE	14	21
Vit D	ug	0.7	1.1
Vit E	mg a-TE	1.25	1.9
Vit K	ug	4.5	6.8
Thiamin	mg	0.15	0.23
Riboflavin	mg mg NE	0.16	0.24 2.3
Niacin	mg NE	1.5	2.3 0.65
Pant. Acid	mg	0.43	0.65
Vit B6	mg	0.15 21	0.23 32
Folic acid Vit B12	ug	070	32 070
	ug	0.30	0.30
Biotin Vit C	ug	4.0 10.0	6.0 15
Choline	mg	28.4	43
Choline Carnitine	mg	28.4 3.0	4.5 4.5
Taurine	mg	10	4.5 15
Idullile	mg	10	10



















DIRECTIONS FOR USAGE

Shake well before opening. Use at room temperature. Collapsible bottles can be attached directly to a Flocare pack giving set. When directly connected to a giving set (aseptically handled, closed system), collapsible bottles have a hanging time of max. 24 hours. If the cap has been removed (used as an open system) then collapsible bottles have a hanging time of max. 8 hours.















Download page






Product leaflet +MIX ONS



Product leaflet +MIX Tube



Product leaflet Tube



Product finder tool